

# Vollständige Literatur zu Christian Dynybil: Rehabilitation nach Hüftendoprothese – was gibt es Neues? OUP 04-2017

## Literatur

- Hudak PL, Hogg-Johnson S, Bombardier C, McKeever PD, Wright JG: Testing a new theory of patient satisfaction with treatment outcome. *Med Care*. 2004; 42: 726–739
- Mahomed NN, Liang MH, Cook EF et al.: The importance of patient expectations in predicting functional outcomes after total joint arthroplasty. *J Rheumatol*. 2002; 29: 1273–1279
- Orbell S, Johnston M, Rowley D, Espley A, Davey P. Cognitive representations of illness and functional and affective adjustment following surgery for osteoarthritis. *Soc Sci Med*. 1998; 471: 93–102
- Kehlet H, Dahl JB: Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet*. 2003; 362: 1921–1928
- Kehlet H, Wilmore D, Multimodal strategies to improve surgical outcome. *Am J Surg*. 2002; 183: 630–641
- Berger R: Podium Präsentation. Rush Medical Center Chicago, IL, AAOS 2016. Orlando/USA
- Dorr LD, Maheshwari AV, Long WT. Early pain relief and function after posterior minimally invasive and conventional total hip arthroplasty. A prospective, randomized, blinded study. *J Bone Joint Surg Am*. 2007; 89: 1153–1160
- Goebel S, Steinert AF, Schillinger J et al.: Reduced postoperative pain in total hip arthroplasty after minimal-invasive anterior approach. *Int Orthop*. 2012; 36: 491498
- Fink B, Mittelstaedt A, Schulz MS, Sebena P, Singer J: Comparison of a minimally invasive posterior approach and the standard posterior approach for total hip arthroplasty. A prospective and comparative study. *J Orthop Surg Res*. 2010; 5: 46
- Mouilhade F, Matsoukis J, Oger P, Mandereau C, Brzakala V, Dujardin F: Component positioning in primary total hip replacement: a prospective comparative study of two anterolateral approaches, minimally invasive versus gluteus medius hemimiotomy. *Orthop Traumatol Surg Res*. 2011; 97: 14–21
- Yang C, Zhu Q, Han Y et al.: Minimally-invasive total hip arthroplasty will improve early postoperative outcomes: a prospective, randomized, controlled trial. *Ir J Med Sci*. 2010; 179: 285–290
- Alecci V, Valente M, Crucil M, Minerva M, Pellegrino CM, Sabbadini DD. Comparison of primary total hip replacements performed with a direct anterior approach versus the standard lateral approach: perioperative findings. *J Orthop Traumatol*. 2011; 12: 123–129
- Jerosch J: Minimalinvasive Hüftendoprothetik. *Dt Ärzteblatt*. 2006; 103: 3333–3339
- Eingartner C. Current trends in total hip arthroplasty. *Ortho Traum Rehab*. 2007; 9: 8–14
- Oganda L, Wilson R, Archbold P et al.: A Minimal-Incision Technique in Total Hip Arthroplasty Does Not Improve Early Postoperative Outcomes. *J Bone Joint Surg Am*. 2005; 87: 701–710
- Spaans AJ, van den Hout JA, Bolder SB: High complication rate in the early experience of minimally invasive total hip arthroplasty by the direct anterior approach. *Acta Orthop*. 2012; 83: 342–346
- Smith TO, Blake V, Hing CB. Minimally invasive versus conventional exposure for total hip arthroplasty: a systematic review and meta-analysis of clinical and radiological outcomes. *Int Orthop*. 2011; 35: 173–184
- Lawlor M, Humphreys P, Morrow E, Ogonda L, Bennett D, Elliott D, Beverland D. Comparison of early postoperative functional levels following total hip replacement using minimally invasive versus standard incisions. A prospective randomized blinded trial. *Clin Rehabil*. 2005; 19: 465–474
- Goebel S, Steinert AF, Schillinger J et al.: Reduced postoperative pain in total hip arthroplasty after minimal-invasive anterior approach. *Int Orthop*. 2012; 36: 491–498
- Peck CN, Foster A, McLauchlan GJ: Reducing incision length or intensifying rehabilitation: what makes the difference to length of stay in total hip replacement in a UK setting? *Int Orthop*. 2006; 30: 395–398
- Pagnano MW, Trousdale RT, Meneghini RM, Hanssen AD: Slower recovery after two-incision than minimally-invasive total hip arthroplasty. A randomized clinical trial. *J Bone Joint Surg Am*. 2008; 90: 1000–1006
- Matta JM, Shahrardar C, Ferguson T: Single incision anterior approach for total hip arthroplasty on an orthopaedic table. *Clin Orthop Relat Res*. 2005; 441: 115–124
- Pagnano MW, Trousdale RT, Meneghini RM, Hanssen AD: Patients preferred a mini-posterior THA to a contralateral two-incision THA. *Clin Orthop Relat Res*. 2006; 453: 156–159
- Dorr LD, Thomas D, Long WT, Polatin PB, Sirianni LE: Psychologic reasons for patients preferring minimally invasive total hip arthroplasty. *Clin Orthop Relat Res*. 2007; 458: 94–100
- Malik A, Dorr LD: The science of minimally invasive total hip arthroplasty. *Clin Orthop Relat Res*. 2007; 463: 74–84
- Camu F, Beecher T, Recker DP, Verburg KM: Valdecoxib, a COX-2-specific inhibitor, is an efficacious, opioid-sparing analgesic in patients undergoing hip arthroplasty. *Am J Ther*. 2002; 9: 43–51
- Husted H, Hansen HC, Holm G et al.: What determines length of stay after total hip and knee arthroplasty? A nationwide study in Denmark. *Arch Orthop Trauma Surg*. 2010; 130: 263–268
- Sharma V, Morgan PM, Cheng EY: Factors influencing early rehabilitation after THA: a systematic review. *Clin Orthop Relat Res*. 2009; 467: 1400–1411
- Horlocker TT, Kopp SL, Pagnano MW, Hebl JR: Analgesia for total hip and knee arthroplasty: a multimodal pathway featuring peripheral nerve block. *J Am Acad Orthop Surg*. 2006; 14: 126–135
- Berger RA, Sanders SA, Thill ES, Sporer SM, Della Valle C. Newer anesthesia and rehabilitation protocols enable outpatient hip replacement in selected patients. *Clin Orthop Relat Res*. 2009; 467:1424–1430
- Long WT, Ward SR, Dorr LD, Raya J, Boutary M, Sirianni LE: Postoperative pain management following total knee arthroplasty: a randomized comparison of continuous epidural versus femoral nerve infusion. *J Knee Surg*. 2006; 19: 137–143
- Maheshwari AV, Boutary M, Yun AG, Sirianni LE, Dorr LD: Multimodal analgesia without routine parenteral narcotics for total hip arthroplasty. *Clin Orthop Relat Res*. 2006; 453: 231–238
- Husted H, Holm G, Jacobsen S: Predictors of length of stay and patient satisfaction after hip and knee replacement surgery: fast-track experience in 712 patients. *Acta Orthop*. 2008; 79: 168–173
- Lunn TH, Husted H, Solgaard S et al.: Intraoperative local infiltration analgesia for early analgesia after total hip arthroplasty: a randomized, double-blind, placebo-controlled trial. *Reg Anesth Pain Med*. 2011; 36: 424–429

35. Parvataneni HK, Shah VP, Howard H, Cole N, Ranawat AS, Ranawat CS: Controlling pain after total hip and knee arthroplasty using a multimodal protocol with local periarticular injections: a prospective randomized study. *J Arthroplasty*. 2007; 22 (6 Suppl 2): 33–38
36. Andersen KV, Pfeiffer-Jensen M, Haraldsted V, Søballe K: Reduced hospital stay and narcotic consumption, and improved mobilization with local and intra-articular infiltration after hip arthroplasty: a randomized clinical trial of an intra-articular technique versus epidural infusion in 80 patients. *Acta Orthop*. 2007; 78: 180–186
37. Linke C, Jaschinski R, Heitmann T et al.: *Biomed* 2013; 1. Berliner Orthopädiereport
38. Mears DC, Mears SC, Chelly JE, Dai F, Vulakovich KL: THA with a minimally invasive technique, multi-modal anesthesia, and home rehabilitation: factors associated with early discharge? *Clin Orthop Relat Res*. 2009; 467: 1412–1417
39. Singelyn FJ, Ferrant T, Malisse MF, Joris D: Effect of intravenous patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous femoral nerve sheath block on rehabilitation after unilateral total-hip arthroplasty. *Reg Anaest Pain Med*. 2005; 30: 452–457
40. Becchi C, Al Malyan M, Coppini R, Campolo M, Magherini M, Boncinelli S: Opioid-free analgesia by continuous psoas compartment block after total hip arthroplasty. A randomized study. *Eur J Anaesthesiol*. 2008; 25: 418–423
41. Sharma V, Morgan PM, Cheng EY: Factors influencing early rehabilitation after THA: a systematic review. *Clin Orthop Relat Res*. 2009; 467: 1400–1411
42. Kehlet H, Slim K. The future of fast-track surgery. *Br J Surg*. 2011; 99: 1025–1026
43. Ilfeld BM, Mariano ER, Girard PJ et al.: A multicenter, randomized, triple-masked, placebo-controlled trial of the effect of ambulatory continuous femoral nerve blocks on discharge-readiness following total knee arthroplasty in patients on general orthopaedic wards. *Pain*. 2010; 150: 477–484
44. McDonald S, Hetrick S, Green S: Pre-operative education for hip or knee replacement. *Cochrane Database Syst Rev*. 2004; (1): CD003526.
45. McDonald S, Page MJ, Behringer K, Wasik J, Sprowson A: Preoperative education for hip or knee replacement. *Cochrane Database Syst Rev*. 2014; (5): CD003526
46. Hozack W: Podium Präsentation, The Rothman Institute, Philadelphia, PA, AAOS 2016. Orlando/USA
47. Minns Lowe CJ, Barker KL, Dewey ME, Sackley CM: Effectiveness of physiotherapy exercise following hip arthroplasty for osteoarthritis: a systematic review of clinical trials. *BMC Musculoskelet Disord*. 2009; 10: 98
48. Minns Lowe CJ, Barker KL, Dewey ME, Sackley CM: Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2007; 335: 812
49. Husted H, Fast-track hip and knee arthroplasty: clinical and organizational aspects. *Acta Orthop Suppl*. 2012; 83: 1–39
50. Peter WF, Nelissen RG, Vlieland TP. Guideline recommendations for post-acute postoperative physiotherapy in total hip and knee arthroplasty: are they used in daily clinical practice? *Musculoskeletal Care*. 2014; 12: 125–31
51. Delorme TL: Restoration of muscle power by heavy-resistance exercises. *J Bone Joint Surg Am*. 1945; 27: 645–667
52. Delorme TL, Watkins AL. Technics of progressive resistance exercise. *Arch Phys Med Rehabil*. 1948 29: 263–273
53. Munin MC, Rudy TE, Glynn NW, Crossett LS, Rubash HE: Early inpatient rehabilitation after elective hip and knee arthroplasty. *JAMA*. 1998; 279: 847–852
54. Bandholm T, Kehlet H: Physiotherapy exercise after fast-track total hip and knee arthroplasty: time for reconsideration? *Arch Phys Med Rehabil*. 2012; 93: 1292–1294
55. Husby VS, Helgerud J, Bjørgen S, Husby OS, Benum P, Hoff J: Early maximal strength training is an efficient treatment for patients operated with total hip arthroplasty. *Arch Phys Med Rehabil*. 2009; 90: 1658–1667
56. Toigo M, Boutellier U: New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur J Appl Physiol*. 2006; 97: 643–663
57. Okoro T, Lemmey AB, Maddison P, Andrew JG: An appraisal of rehabilitation regimes used for improving functional outcome after total hip replacement surgery. *Sports Med Arthrosc Rehabil Ther Technol*. 2012; 4–5
58. Lenssen AF, Crijns YH, Waltjé EM et al.: Efficiency of immediate postoperative inpatient physical therapy following total knee arthroplasty: an RCT. *BMC Musculoskelet Disord*. 2006; 7: 71
59. Moffet H, Collet JP, Shapiro SH, Paradis G, Marquis F, Roy L: Effectiveness of intensive rehabilitation on functional ability and quality of life after first total knee arthroplasty: A single-blind randomized controlled trial. *Arch Phys Med Rehabil*. 2004; 85: 546–556
60. Brower RG, Consequences of bed rest. *Crit Care Med*. 2009; 37(10 Suppl): 422–428
61. Piva SR, Teixeira PE, Almeida GJ et al.: Contribution of hip abductor strength to physical function in patients with total knee arthroplasty. *Phys Ther*. 2011; 91: 225–233
62. Toigo M, Boutellier U: New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur J Appl Physiol*. 2006; 97: 643–663
63. Silkman Baker C, Mc Keon JM: Does pre-operative rehabilitation improve patient – based outcomes in persons who have undergone total knee arthroplasty? A systematic review. *PM R* 2012; 4: 756–767
64. Lenssen AF, de Bie RA: Role of physiotherapy in peri-operative management in total knee and hip surgery. *Injury*. 2006; 37 Suppl 5: 41–43
65. Khan F, Ng L, Gonzalez S, Hale T, Turner-Stokes L: Multidisciplinary rehabilitation programmes following joint replacement at the hip and knee in chronic arthropathy. *Cochrane Database Syst Rev*. 2008; (2): CD004957
66. Tayrose G, Newman D, Slover J, Jaffe F, Hunter T, Bosco J 3rd: Rapid mobilization decreases length-of-stay in joint replacement patients. *Bull Hosp Jt Dis*. 2013; 71: 222–226
67. Husby VS, Helgerud J, Bjørgen S, Husby OS, Benum P, Hoff J: Early postoperative maximal strength training improves work efficiency 6–12 months after osteoarthritis-induced total hip arthroplasty in patients younger than 60 years. *Am J Phys Med Rehabil* 2010; 89: 304–314
68. Kristensen J, Franklyn-Miller A. Resistance training in musculoskeletal rehabilitation: a systematic review. *Br J Sports Med*. 2012; 46: 719–726
69. Krank aus dem Krankenhaus. Visite. [www.ndr.de/Ratgeber/Gesundheit/Zufrueh-entlassen-krank-aus-dem-krankenhaus](http://www.ndr.de/Ratgeber/Gesundheit/Zufrueh-entlassen-krank-aus-dem-krankenhaus). Tina Roth. 26.05.2015
70. Greitemann B: REDIA-Studie. *Die Welt*, 2011
71. Heisel J, Jerosch J: Rehabilitationsfähigkeit – Rehabilitation nach Hüft- und Knieendoprothese. Köln: Deutscher Ärzte-Verlag, 2007: 71–73
72. Linke C, Jaschinski T, Heitmann T et al.: 1. Berliner Orthopädiereport. *Biomet*. 2013
73. Diehl R, Gebauer E, Groner A: *Kursbuch Sozialmedizin. Lehrbuch zum Curriculum der Bundesärztekammer*. Köln: Deutscher Ärzte-Verlag, 2012
74. Eiff W, Schüring S, Niehus C: REDIA. Auswirkungen der DRG – Einführung Rehabilitation. Berlin: Lit-Verlag, 2011
75. Statistisches Bundesamt. Grunddaten der Vorsorge- oder Rehabilitationseinrichtungen. Wiesbaden, 2009
76. Eiff W: Krankenhaus-IT Journal. Forschung. Ausgabe 03/2011
77. Reha-Bericht 2015, 6. Auflage, Deutsche Rentenversicherung Bund
78. Steiner M, Zwinkmann C, Riedel W et al.: Die medizinische Rehabilitation Erwerbstätiger – Sicherung von Produktivität und Wachstum. Prognos AG 2009
79. Eiff W, Schüring S: Kürzere Akutverweildauern erhöhen Aufwand in der Reha. *Dt. Ärzteblatt* 2011; 108: 1164–1166