

Vollständige Literatur zu Klaus-Peter Günther et al.: Wann ist der richtige Zeitpunkt für die Primäroperation in der Hüftendoprothetik? OUP 04-2017

Literatur

1. Balck F, Lippmann M, Jeszenszky C, Günther KP, Kirschner S: The Influence of Optimism on Functionality after Total Hip Replacement Surgery. *J Health Psychol* 2015 Jan 20. pii: 1359105314566256. [Epub ahead of print]
2. Bleß H, Kip M (Hrsg.): Weißbuch Gelekersatz. Versorgungssituation endoprothetischer Hüft- und Knieoperationen in Deutschland. Springer 2016
3. Bozic KJ, Lau E, Kurtz S et al.: Patient-related risk factors for periprosthetic joint infection and postoperative mortality following total hip arthroplasty in Medicare patients. *J Bone Joint Surg Am* 2012; 94: 794–800
4. Bozic KJ, Lau E, Ong K et al.: Risk factors for early revision after primary total hip arthroplasty in Medicare patients. *Clin Orthop Relat Res* 2014; 472: 449–54
5. Clement ND, Muzammil A, Macdonald D, Howie CR, Biant LC: Socioeconomic status affects the early outcome of total hip replacement. *The Journal of bone and joint surgery* 2011; British volume 93: 464–69
6. Deutsches Institut für Medizinische Dokumentation und Information (DIMDI) (Hrsg.) Internationale Klassifikation der Funktionsfähigkeit, Behinderung und Gesundheit (ICF), Genf: WHO, 2005
7. Dreinhöfer KE, Dieppe P, Stürmer T et al.: Indications for total hip replacement: comparison of assessments of orthopaedic surgeons and referring physician. *Ann Rheum Dis* 2006; 65: 1346–50
8. Dowsey MM, Nikpour M, Choong PF: 2014 Outcomes following large joint arthroplasty: does socio-economic status matter? *BMC Musculoskelet Disor* 2014; 15: 148
9. Finkenstädt V, Niehaus F: Die Aus sagekraft von Länderrankings im Ge sundheitsbereich. Köln: Wissenschaftliches Institut der PKV, 2015
10. Fogel GR, Esses SI: Hip spine syndrome: management of coexisting radiculopathy and arthritis of the lower extremity. *The Spine Journal* 2003; 3: 238–41
11. Gossec L, Paternotte S, Maillefert JF: et al: The role of pain and functional impairment in the decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1909 patients. Report of the OARSI-OMERACT Task Force on total joint replacement. *Osteoarthritis Cartilage* 2011; 19: 147–54
12. Greene ME, Rolfson O, Nemes S, Gordon M, Malchau H, Garellick G: Education attainment is associated with patient-reported outcomes: findings from the Swedish Hip Arthroplasty Register. *Clin Orthop Relat Res* 2014; 472: 1868–76
13. Günther KP, Haase E, Lange T et al.: Persönlichkeitsprofil und Komorbidität: Gibt es den „schwierigen Patienten“ in der primären Hüftendoprothetik? *Orthopäde* 2015; 44: 555–65
14. Haanstra TM, van den Berg T, Ostelo RW et al.: Systematic review: do patient expectations influence treatment outcomes in total knee and total hip arthroplasty? *Health Qual Life Outcomes* 2012; 10: 152
15. Haase E, Kopkow C, Beyer F et al.: Patient-reported outcomes and outcome predictors after primary total hip arthroplasty: results from the Dresden Hip Registry. *Hip Int* 2015; Oct 6:0. doi: 10.5301/hipint.5000300. [Epub ahead of print]
16. Hamann C, Günther KP, Lützner J, Beyer F, Kirschner S: Diabetes mellitus – Ein unterschätzter Risikofaktor in der Hüftendoprothetik? *Osteologie* 2015; (article in press)
17. Hamilton DF, Lane JV, Gaston P et al.: What determines patient satisfaction with surgery? A prospective cohort study of 4709 patients following total joint replacement. *BMJ Open* 2013; 3(4). pii: e002525. doi: 10.1136/bmjjopen-2012-002525. Print 2013
18. Haverkamp D, Klinkenbijl MN, Somford MP, Albers GH, van der Vis HM: Obesity in total hip arthroplasty – does it really matter? A meta-analysis. *Acta Orthop* 2011; 82: 417–22
19. Hawker GA, Badley EM, Borkhoff CM et al.: Which patients are most likely to benefit from total joint arthroplasty? *Arthritis Rheum* 2013; 65: 1243–52
20. Herberth P, Malchau H: Long-term registration has improved the quality of hip replacement: A review of the Swedish THR Register comparing 160,000 cases. *Acta Orthop* 2000; 71: 111–21
21. Hooper GJ, Rothwell AG, Hooper NM, Frampton C: The relationship between the American Society Of Anesthesiologists physical rating and outcome following total hip and knee arthroplasty: an analysis of the New Zealand Joint Registry. *J Bone Joint Surg Am* 2012; 94: 1065–70
22. Hutchings A, Neuburger J, Grosse Frie K, Black N, van der Meulen J: Factors associated with non-response in routine use of patient reported outcome measures after elective surgery in England. *Health Qual Life Outcomes* 2012; 10: 34
23. Judge A, Arden NK, Batra RN, Thomas G, Beard D, Javaid MK, et al.: The association of patient characteristics and surgical variables on symptoms of pain and function over 5 years following primary hip-replacement surgery: a prospective cohort study. *BMJ Open* 2013; 3: e002453
24. Judge A, Cooper C, Williams S, Dreinhöfer K, Dieppe P: Patient-reported outcomes one year after primary hip replacement in a European Collaborative Cohort. *Arthritis Care Res* 2010; 62: 480–488
25. Judge A, Cooper C, Arden NK et al.: Pre-operative expectation predicts 12-month post-operative outcome among patients undergoing primary total hip replacement in European orthopaedic centres. *Osteoarthritis Cartilage* 2011; 19: 659–67
26. Judge A, Javaid MK, Arden NK et al.: Clinical tool to identify patients who are most likely to achieve long-term improvement in physical function after total hip arthroplasty. *Arthritis Care Res* 2012; 64: 881–9
27. Judge A, Batra RN, Thomas GE et al.: Body mass index is not a clinically meaningful predictor of patient reported outcomes of primary hip replacement surgery: prospective cohort study. *Osteoarthritis Cartilage* 2014; 22: 431–9
28. Jüni P, Low N, Reichenbach S, Villiger PM, Williams S, Dieppe PA: Gender inequity in the provision of care for hip disease: population-based cross-sectional study. *Osteoarthritis Cartilage* 2010; 18: 640–5
29. Kapoor A, Chew P, Silliman RA et al.: Venous thromboembolism after joint replacement in older male veterans with comorbidity. *J Am Geriatr Soc* 2013; 61: 590–601
30. Mancuso CA, Jout J, Salvati EA, Sculco TP: Fulfillment of patients' expectations for total hip arthroplasty. *The Journal of bone and joint surgery* 2009; American volume 91: 2073–78

31. Martinez-Huedo MA, Villanueva M, de Andres AL et al.: Trends 2001 to 2008 in incidence and immediate postoperative outcomes for major joint replacement among Spanish adults suffering diabetes. *Eur J Orthop Surg Traumatol* 2013; 23: 53–9
32. Neuburger J, Hutchings A, Black N, van der Meulen JH: Socioeconomic differences in patient-reported outcomes after a hip or knee replacement in the English National Health Service. *Journal of Public Health* 2013; 35: 115–24
33. Nilssdotter A: Predictors of patient relevant outcome after total hip replacement for osteoarthritis: a prospective study. *Annals of the Rheumatic Diseases* 2003; 62: 923–30
34. Palazzo C, Jourdan C, Descamps S et al.: Determinants of satisfaction 1 year after total hip arthroplasty: the role of expectations fulfilment. *BMC Musculoskeletal Disorders* 2014; 15: 53
35. Peters ML, Sommer M, de Rijke JM et al.: Somatic and psychologic predictors of long-term unfavorable outcome after surgical intervention. *Ann Surg* 2007; 245: 487–94
36. Prokopetz JJ, Losina E, Bliss RL, Wright J, Baron JA, Katz JN: Risk factors for revision of primary total hip arthroplasty: a systematic review. *BMC Musculoskeletal Disorders* 2012; 13: 251
37. Quintana JM, Escobar A, Aguirre U, Lafuente I, Arenaza JC: Predictors of Health-related Quality-of-life Change after Total Hip Arthroplasty. *Clin Orthop Relat Res* 2009; 467: 2886–94
38. Rolfson O, Dahlberg LE, Nilsson JA et al.: Variables determining outcome in total hip replacement surgery. *J Bone Joint Surg Br* 2009; 91: 157–61
39. Santaguida PL, Hawker GA, Hudak PL et al.: Patient characteristics affecting the prognosis of total hip and knee joint arthroplasty: a systematic review. *Can J Surg* 2008; 51: 428–36
40. Schäfer T, Pritzkuleit R, Jeszenszky C et al.: Trends and geographical variation of primary hip and knee joint replacement in Germany. *Osteoarthritis and Cartilage* 2013; 21: 279–88
41. Schäfer T, Krummenauer F, Mettelsiefen J, Kirschner S, Günther KP: Social, educational, and occupational predictors of total hip replacement outcome. *Osteoarthritis and Cartilage* 2010; 18: 1036–42
42. Schäfer T, Biakowski S, Walther A, Fickert S, Krummenauer F, Günther KP: Preoperative importance of improvement and postoperative satisfaction in patients undergoing total hip replacement. *Eur Orthop Traumatol* 2011; 1: 143–52
43. Singh JA, Lewallen DG: Ipsilateral lower extremity joint involvement increases the risk of poor pain and function outcome after hip or knee arthroplasty. *BMC Med* 2013; 11: 144
44. Stiehler M, Goronzy J, Günther KP: Die endoprothetische Versorgung beim übergewichtigen Koxarthrosepatienten. *Orthopäde* 2015; 44: 523–30
45. Stürmer T, Dreinhöfer K, Gröber-Grätz D et al.: Differences in the views of orthopaedic surgeons and referring practitioners on the determinants of outcome after total hip replacement. *J Bone Joint Surg Br* 2005; 87: 1416–9
46. Tsang ST, Gaston P: Adverse peri-operative outcomes following elective total hip replacement in diabetes mellitus: a systematic review and meta-analysis of cohort studies. *Bone Joint J* 2013; 95-B: 1474–9
47. Vissers MM, Bussmann JB, Verhaar JA et al.: Psychological Factors Affecting the Outcome of Total Hip and Knee Arthroplasty: A Systematic Review. *Semin Arthritis Rheum* 2012; 41: 576–588
48. Young NL, Cheah D, Waddell JP, Wright JG: Patient characteristics that affect the outcome of total hip arthroplasty: a review. *Can J Surg* 1998; 41: 188–95
49. Zmistrovski B1, Dizdarevic I, Jacovides CL, Radcliff KE, Mraovic B, Parvizi J: Patients with uncontrolled components of metabolic syndrome have increased risk of complications following total joint arthroplasty. *J Arthroplasty* 2013; 28: 904–7