

# Vollständige Literatur zu Carsten Tibesku: Gelenkersetzende Therapie bei Gonarthrose OUP 07-2018

## Literatur

1. DGOU DGfOuU. Leitlinie Gonarthrose. [www.awmf.org/leitlinien/detail/ll/033-004.html](http://www.awmf.org/leitlinien/detail/ll/033-004.html) 2018
2. Berchuck M, Andriacchi TP, Bach BR, Reider B: Gait adaptations by patients who have a deficient anterior cruciate ligament. *J Bone Joint Surg Am.* 1990; 72: 871–7
3. Andriacchi TP: Dynamics of pathological motion: applied to the anterior cruciate deficient knee. *J Biomech.* 1990; 23 Suppl 1: 99–105
4. Chassin EP, Mikosz RP, Andriacchi TP, Rosenberg AG: Functional analysis of cemented medial unicompartmental knee arthroplasty. *J Arthroplasty.* 1996; 11: 553–9
5. Argenson JN, Komistek RD, Aubaniac JM et al.: In vivo determination of knee kinematics for subjects implanted with a unicompartmental arthroplasty. *J Arthroplasty.* 2002; 17: 1049–54
6. Becker R, Mauer C, Starke C et al.: Anteroposterior and rotational stability in fixed and mobile bearing unicondylar knee arthroplasty: a cadaveric study using the robotic force sensor system. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21: 2427–32
7. Fuchs S, Tibesku CO, Flören M, Thorwesken L: Interdependence of clinical and isokinetic results after bicondylar knee prostheses with special emphasis on quality of life results. *Int Orthop.* 2000; 24: 268–71
8. Fuchs S, Tibesku CO, Genkinger M, Laass H, Rosenbaum D: Proprioception with bicondylar sledge prostheses retaining cruciate ligaments. *Clin Orthop Relat Res.* 2003; 148–54
9. Fuchs S, Tibesku CO, Frisse D, Laass H, Rosenbaum D: Quality of life and gait after unicondylar knee prosthesis are inferior to age-matched control subjects. *Am J Phys Med Rehabil.* 2003; 82: 441–6
10. Dalury DF, Fisher DA, Adams MJ, Gonzales RA: Unicompartmental knee arthroplasty compares favorably to total knee arthroplasty in the same patient. *Orthopedics.* 2009; 32 (4)
11. Von Keudell A, Sodha S, Collins J, Minas T, Fitz W, Gomoll AH: Patient satisfaction after primary total and unicompartmental knee arthroplasty: an age-dependent analysis. *The Knee.* 2014; 21: 180–4
12. Berend KR, Berend ME, Dalury DF, Argenson JN, Dodd CA, Scott RD: Consensus Statement on Indications and Contraindications for Medial Unicompartimental Knee Arthroplasty. *J Surg Orthop Adv.* 2015; 24: 252–6
13. Niinimaki TT, Murray DW, Partanen J, Pajala A, Leppilahti JI: Unicompartmental knee arthroplasties implanted for osteoarthritis with partial loss of joint space have high re-operation rates. *The Knee.* 2011; 18: 432–5
14. Pandit H, Gulati A, Jenkins C et al.: Unicompartmental knee replacement for patients with partial thickness cartilage loss in the affected compartment. *The Knee.* 2011; 18: 168–71
15. Cheng M, Chen D, Guo Y, Zhu C, Zhang X: Comparison of fixed- and mobile-bearing total knee arthroplasty with a mean five-year follow-up: A meta-analysis. *Experimental and therapeutic medicine.* 2013; 6: 45–51
16. Kleebald LJ, van der List JP, Zuiderbaan HA, Pearle AD: Larger range of motion and increased return to activity, but higher revision rates following unicompartmental versus total knee arthroplasty in patients under 65: a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2017 Nov 28 (Epub ahead of print)
17. Svärd UC, Price AJ: Oxford medial unicompartmental knee arthroplasty. A survival analysis of an independent series. *J Bone Joint Surg Br.* 2001; 83: 191–4
18. Heyse TJ, Tibesku CO: Lateral unicompartmental knee arthroplasty: a review. *Arch Orthop Trauma Surg.* 2010; 130: 1539–48
19. Volpi P, Marinoni L, Bait C, Galli M, Denti M: Lateral unicompartmental knee arthroplasty: indications, technique and short-medium term results. *Knee Surg Sports Traumatol Arthrosc.* 2007; 15: 1028–34
20. Walker T, Zahn N, Bruckner T et al.: Mid-term results of lateral unicondylar mobile bearing knee arthroplasty: a multicentre study of 363 cases. *Bone Joint J.* 2018; 100-B(1): 42–49
21. Argenson JN, Parratte S, Ashour A, Komistek RD, Scuderi GR: Patient-reported outcome correlates with knee function after a single-design mobile-bearing TKA. *Clin Orthop Relat Res.* 2008; 466: 2669–76
22. Pisanu G, Rosso F, Bertolo C et al.: Patellofemoral Arthroplasty: Current Concepts and Review of the Literature. *Joints.* 2017; 5: 237–45
23. Leadbetter WB, Ragland PS, Mont MA: The appropriate use of patellofemoral arthroplasty: an analysis of reported indications, contraindications, and failures. *Clin Orthop Relat Res.* 2005 (436): 91–9
24. Beckmann J, Steinert A, Zilkens C et al.: [Partial replacement of the knee joint with patient-specific instruments and implants (ConforMIS iUni, iDuo)]. *Der Orthopäde.* 2016; 45: 322–30
25. Parratte S, Ollivier M, Opsomer G, Lunebourg A, Argenson JN, Thienpont E: Is knee function better with contemporary modular bicompartamental arthroplasty compared to total knee arthroplasty? Short-term outcomes of a prospective matched study including 68 cases. *Orthop Traumatol Surg Res.* 2015; 101: 547–52
26. Benazzo F, Rossi SM, Ghira M: Partial knee arthroplasty: patellofemoral arthroplasty and combined unicompartmental and patellofemoral arthroplasty implants-general considerations and indications, technique and clinical experience. *The Knee.* 2014; 21 Suppl 1: S43–6
27. Thienpont E, Price A: Bicompartamental knee arthroplasty of the patellofemoral and medial compartments. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21: 2523–31
28. Yeo NE, Chen JY, Yew A, Chia SL, Lo NN, Yeo SJ: Prospective randomised trial comparing unlinked, modular bicompartamental knee arthroplasty and total knee arthroplasty: a five years follow-up. *The Knee.* 2015; 22: 321–7
29. Fuchs S, Tibesku CO, Frisse D, Genkinger M, Laass H, Rosenbaum D: Clinical and functional comparison of uni- and bicondylar sledge prostheses. *Knee Surg Sports Traumatol Arthrosc.* 2005; 13: 197–202
30. Gunston FH: Polycentric knee arthroplasty. Prosthetic simulation of normal knee movement. *J Bone Joint Surg Br.* 1971; 53: 272–7
31. Coventry MB, Finerman GA, Riley LH, Turner RH, Upshaw JE: A new geometric knee for total knee arthroplasty. *Clin Orthop Relat Res.* 1972; 83: 157–62
32. Riley D, Woodyard JE: Long-term results of Geomedic total knee replacement. *J Bone Joint Surg Br.* 1985; 67: 548–50
33. Cloutier JM: Results of total knee arthroplasty with a non-constrained prosthesis. *J Bone Joint Surg Am.* 1983; 65: 906–19
34. Cloutier JM, Sabouret P, Deghrar A: Total knee arthroplasty with retention of

- both cruciate ligaments. A nine to eleven-year follow-up study. *J Bone Joint Surg Am.* 1999; 81: 697–702
35. Baumann F, Krutsch W et al.: Reduced joint-awareness in bicruciate-retaining total knee arthroplasty compared to cruciate-sacrificing total knee arthroplasty. *Arch Orthop Trauma Surg.* 2018; 138: 273–9
36. Baumann F, Bahadin O, Krutsch W et al.: Proprioception after bicruciate-retaining total knee arthroplasty is comparable to unicompartmental knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2017; 25:1697–704
37. Athwal KK, Hunt NC, Davies AJ, Deehan DJ, Amis AA: Clinical biomechanics of instability related to total knee arthroplasty. *Clin Biomech (Bristol, Avon).* 2014; 29: 119–28
38. Ishii Y, Noguchi H, Takeda M, Kiga H, Toyabe SI: Effect of voluntary soft tissue tension and articular conformity after total knee arthroplasty on in vivo anteroposterior displacement. *The Knee.* 2011; 18: 11–4
39. Haider H, Walker PS: Measurements of constraint of total knee replacement. *J Biomech.* 2005; 38: 341–8
40. Heesterbeek P, Keijser N, Jacobs W, Verdonschot N, Wymenga A: Posterior cruciate ligament recruitment affects antero-posterior translation during flexion gap distraction in total knee replacement. An intraoperative study involving 50 patients. *Acta orthopaedica.* 2010; 81: 471–7
41. Zelle J, Heesterbeek PJ, De Waal Malefijt M, Verdonschot N: Numerical analysis of variations in posterior cruciate ligament properties and balancing techniques on total knee arthroplasty loading. *Med Eng Phys.* 2010; 32: 700–7
42. Fantozzi S, Catani F, Ensini A, Leardini A, Giannini S: Femoral rollback of cruciate-retaining and posterior-stabilized total knee replacements: in vivo fluoroscopic analysis during activities of daily living. *J Orthop Res.* 2006; 24: 2222–9
43. Walker PS, Sussman-Fort JM, Yildirim G, Boyer J: Design features of total knees for achieving normal knee motion characteristics. *J Arthroplasty.* 2009; 24: 475–83
44. Lattanzio PJ, Chess DG, MacDermid JC: Effect of the posterior cruciate ligament in knee-joint proprioception in total knee arthroplasty. *J Arthroplasty.* 1998; 13: 580–5
45. Matziolis G, Mehlhorn S, Schattat N et al.: How much of the PCL is really preserved during the tibial cut? *Knee Surg Sports Traumatol Arthrosc.* 2012; 20: 1083–6
46. Moro-oka TA, Muenchinger M, Canciani JP, Banks SA: Comparing in vivo kinematics of anterior cruciate-retaining and posterior cruciate-retaining
- total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2007; 15: 93–9
47. Victor J, Banks S, Bellemans J: Kinematics of posterior cruciate ligament-retaining and -substituting total knee arthroplasty: a prospective randomised outcome study. *J Bone Joint Surg Br.* 2005; 87: 646–55
48. Bellemans J, Banks S, Victor J, Vandenneucker H, Moemans A: Fluoroscopic analysis of the kinematics of deep flexion in total knee arthroplasty. Influence of posterior condylar offset. *J Bone Joint Surg Br.* 2002; 84: 50–3
49. Watanabe T, Ishizuki M, Muneta T, Banks SA: Knee kinematics in anterior cruciate ligament-substituting arthroplasty with or without the posterior cruciate ligament. *J Arthroplasty.* 2013; 28: 548–52
50. Heyse TJ, Becher C, Kron N et al.: Quadriceps force in relation of intrinsic anteroposterior stability of TKA design. *Arch Orthop Trauma Surg.* 2010; 130: 1–9
51. Becher C, Heyse TJ, Kron N et al.: Posterior stabilized TKA reduce patellofemoral contact pressure compared with cruciate retaining TKA in vitro. *Knee Surg Sports Traumatol Arthrosc.* 2009; 17: 1159–65
52. Bellemans J, Robijns F, Duerinckx J, Banks S, Vandenneucker H: The influence of tibial slope on maximal flexion after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2005; 13: 193–6
53. Arabori M, Matsui N, Kuroda R et al.: Posterior condylar offset and flexion in posterior cruciate-retaining and posterior stabilized TKA. *J Orthop Sci.* 2008; 13: 46–50
54. Jacobs WC, Clement DJ, Wymenga AB: Retention versus removal of the posterior cruciate ligament in total knee replacement: a systematic literature review within the Cochrane framework. *Acta orthopaedica.* 2005; 76: 757–68
55. Abdel MP, Morrey ME, Jensen MR, Morrey BF: Increased long-term survival of posterior cruciate-retaining versus posterior cruciate-stabilizing total knee replacements. *J Bone Joint Surg Am.* 2011; 93: 2072–8
56. National Joint Replacement Registry, annual report [database on the Internet]. 2011 [cited 2013]
57. Bercik MJ, Joshi A, Parvizi J: Posterior cruciate-retaining versus posterior-stabilized total knee arthroplasty: a meta-analysis. *J Arthroplasty.* 2013; 28: 439–44
58. Lombardi AV, Jr., Berend KR: Posterior cruciate ligament-retaining, posterior stabilized, and varus/valgus posterior stabilized constrained articulations in total knee arthroplasty. *Instr Course Lect.* 2006; 55: 419–27
59. Laskin RS, Maruyama Y, Villaneuva M, Bourne R: Deep-dish congruent tibial component use in total knee arthroplasty: a randomized prospective study. *Clin Orthop Relat Res.* 2000; 380: 36–44
60. Scott RD, Thornhill TS: Posterior cruciate supplementing total knee replacement using conforming inserts and cruciate recession. Effect on range of motion and radiolucent lines. *Clin Orthop Relat Res.* 1994; 309: 146–9
61. Daniilidis K, Skwara A, Vieth V et al.: Highly conforming polyethylene inlays reduce the in vivo variability of knee joint kinematics after total knee arthroplasty. *The Knee.* 2012; 19: 260–5
62. Fisher J, Jennings LM, Galvin AL, Jin ZM, Stone MH, Ingham E: 2009 Knee Society Presidential Guest Lecture: Polyethylene wear in total knees. *Clin Orthop Relat Res.* 2010; 468: 12–8
63. Galvin AL, Kang L, Udofoia I et al.: Effect of conformity and contact stress on wear in fixed-bearing total knee prostheses. *J Biomech.* 2009; 42: 1898–902
64. Huang CH, Ma HM, Liau JJ, Ho FY, Cheng CK: Osteolysis in failed total knee arthroplasty: a comparison of mobile-bearing and fixed-bearing knees. *J Bone Joint Surg Am.* 2002; 84-A (12): 2224–9
65. Pagnano MW, Trousdale RT, Stuart MJ, Hanssen AD, Jacofsky DJ: Rotating platform knees did not improve patellar tracking: a prospective, randomized study of 240 primary total knee arthroplasties. *Clin Orthop Relat Res.* 2004; 428: 221–7
66. Jacobs WC, Clement DJ, Wymenga AB: Retention versus sacrifice of the posterior cruciate ligament in total knee replacement for treatment of osteoarthritis and rheumatoid arthritis. *Cochrane Database Syst Rev.* 2005 (4): CD004803.
67. van der Voort P, Pijls BG, Nouta KA, Valstar ER, Jacobs WC, Nelissen RG: A systematic review and meta-regression of mobile-bearing versus fixed-bearing total knee replacement in 41 studies. *Bone Joint J.* 2013; 95-B (9): 1209–16
68. Girard J, Amzallag M, Pasquier G et al.: Total knee arthroplasty in valgus knees: predictive preoperative parameters influencing a constrained design selection. *Orthop Traumatol Surg Res.* 2009; 95: 260–6
69. Sculco TP: The role of constraint in total knee arthroplasty. *J Arthroplasty.* 2006; 21 (4 Suppl 1): 54–6
70. Gustke KA: Preoperative planning for revision total knee arthroplasty: avoiding chaos. *J Arthroplasty.* 2005; 20 (4 Suppl 2): 37–40
71. McAuley JP, Engh GA: Constraint in total knee arthroplasty: when and what? *J Arthroplasty.* 2003; 18 (3 Suppl 1): 51–4
72. Bauwens K, Matthes G, Wich M et al.: Navigated total knee replacement. A meta-analysis. *J Bone Joint Surg Am.* 2007; 89: 261–9

73. Tibesku CO: [Total knee arthroplasty with the use of patient specific instruments. The VISIONAIRE system]. Der Orthopäde. 2016; 45: 286–93
74. Mannan A, Akinyooye D, Hossain F: A Meta-analysis of Functional Outcomes in Patient-Specific Instrumented Knee Arthroplasty. Knee Surg. 2017; 30: 668–74
75. Mannan A, Smith TO: Favourable rotational alignment outcomes in PSI knee arthroplasty: A Level 1 systematic review and meta-analysis. The Knee. 2016; 23: 186–90
76. Tibesku CO, Hofer P, Portegies W, Ruys CJ, Fennema P: Benefits of using customized instrumentation in total knee arthroplasty: results from an activity-based costing model. Arch Orthop Trauma Surg. 2013; 133: 405–11
77. Slamin J, Parsley B: Evolution of customization design for total knee arthroplasty. Curr Rev Musculoskelet Med. 2012; 5: 290–5
78. Steinert AF, Holzapfel BM, Sefrin L, Arnholdt J, Hoberg M, Rudert M: Tota-lendoprothetischer Kniegelenkersatz. Patientenspezifische Instrumente und Implantate. Der Orthopäde. 2016; 45: 331–40
79. Jerosch J, Heisel J, Tibesku CO: Knieendoprothetik, Indikationen, Operationstechnik, Nachbehandlung, Begutachtung, Berlin: Springer-Verlag, 2. Aufl. 2015